

PRODUCERS

303/10695

STOCKER PRO FEEDLOT R20 DFBZ

MEDICATED

FOR GROWING CATTLE ON PASTURE OR IN DRYLOT

For increased rate of weight gain. An insect growth regulator, which prevents the development of house, stable, face and horn flies in the manure of treated beef and dairy cattle.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium)	20.0 g/ton
Diflubenzuron.....	6.05 g/ton

GUARANTEED ANALYSIS

Crude Protein	(Min)	13.00 %	Phosphorus	(Min)	0.50 %
Crude Fat	(Min)	3.75 %	Salt	(Min)	0.30 %
Crude Fiber	(Max)	11.00 %	Salt	(Max)	0.70 %
Calcium	(Min)	0.65 %	Potassium	(Min)	0.75 %
Calcium	(Max)	1.05 %	Vitamin A	(Min)	8,000 IU/LB

INGREDIENTS

Grain Products, Plant Protein Products, Roughage Products (13%), Processed Grain Byproducts, Molasses Products, Ground Limestone, Salt, Monocalcium Phosphate, Calcium Propionate, Selenium Yeast, Cobalt Carbonate, Copper Chloride, Copper Sulfate, Ethylenediamine Dihydriodide, Manganese Sulfate, Manganous Oxide, Zinc Sulfate, Zinc Oxide, Mineral Oil, Vitamin E Supplement, Vitamin A Supplement and Vitamin D3 Supplement.

FEEDING DIRECTIONS

Feed this product at a rate of 5 to 20 pounds per head per day to provide 50 to 200 milligrams monensin per head per day and 0.1 mg diflubenzuron/kg of bodyweight. Always provide plenty of fresh, clean water.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. Must be thoroughly mixed in feeds before use. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

CAUTION: Keep out of reach of children. Discard empty container according to local regulations. Never reuse empty container.

WARNING: A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

Manufactured by
PRODUCERS COOPERATIVE ASSOCIATION
Bryan, Texas 77806

Net Weight 50 lbs. (22.68 kg) or Bulk