

PRODUCERS

44376/13190

CATTLE DEWORMER CUBE

MEDICATED

FOR CATTLE ON PASTURE

For the removal and control of lungworms (*Dictyocaulus viviparus*) barberpole worms (*Haemonchus contortus*); brown stomach worms (*Ostertagia ostertagi*); small stomach worms (*Trichostrongylus axei*); hookworms (*Bunostomum phlebotomum*); thread-necked intestinal worms (*Nematodirus helvetianus*); small intestinal worms (*Cooperia punctatata* and *Cooperia oncophora*); bankrupt worms (*Trichostrongylus colubriformis*); and nodular worms (*Oesophagostomum radiatum*).

ACTIVE DRUG INGREDIENT

Fenbendazole 908 g/ton

GUARANTEED ANALYSIS

Crude Protein	(Min)	20.00 %	Phosphorus	(Min)	0.75 %
Crude Fat	(Min)	2.60 %	Salt	(Min)	0.75 %
Crude Fiber	(Max)	10.00 %	Salt	(Max)	1.25 %
Calcium	(Min)	0.75 %	Potassium	(Min)	1.00 %
Calcium	(Max)	1.25 %	Vitamin A	(Min)	8,700 IU/LB

INGREDIENTS

Processed Grain Byproducts, Cottonseed Meal, Grain Products, Molasses Products, Ground Limestone, Salt, Calcium Lignosulfonate, Vitamin A Supplement, Potassium Chloride, Calcium Propionate, Selenium Yeast, Cobalt Carbonate, Copper Chloride, Copper Sulfate, Ethylenediamine Dihydriodide, Manganese Sulfate, Manganous Oxide, Zinc Sulfate, Zinc Oxide, Mineral Oil and Calcium Carbonate.

FEEDING DIRECTIONS

Feed as the sole ration for one day. Feed at the rate of 0.5 pound of Cattle Dewormer Cubes per 100 pounds of bodyweight to provide 2.27 milligrams fenbendazole per pound of bodyweight. See the table for additional feeding amounts based on various beef cattle bodyweights. Sufficient space must be available so all animals can eat at the same time to ensure uniform consumption.

Bodyweight	Amt. Deworm. Cube/Day
250	1.25
500	2.5
750	3.75
1,000	5.0
1250	6.25

CAUTION: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

LIMITATIONS: Feed as sole ration or top dress for one day. Retreatment may be needed after 4 to 6 weeks. Cattle must not be slaughtered within 13 days following last treatment. Do not use within 13 days of slaughter. For dairy cattle the milk discard is zero hours.

Note: Consult veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Manufactured by
PRODUCERS COOPERATIVE ASSOCIATION
Bryan, Texas 77806

Net Weight 50 lbs. (22.68 kg) or Bulk

11114 67 5/5/2020

MFG: ~DATE ~MTIME

