

PRODUCERS

478/992 SHOW GOAT

MEDICATED FOR GROWING GOATS (FOR RUMINANTS ONLY)

For the prevention of coccidiosis caused by *Eimeria crandalis*, *Eimeria christenseni* and *Eimeria ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT
Monensin (as monensin sodium) 20 g/ton

GUARANTEED ANALYSIS

Crude Protein	(Min)	16.00 %			
Includes not more than 1.15% equivalent non-protein nitrogen.					
Crude Fat	(Min)	3.70 %	Salt	(Min)	0.80 %
Crude Fiber	(Max)	14.00 %	Salt	(Max)	1.20 %
Calcium	(Min)	0.80 %	Copper	(Min)	29 PPM
Calcium	(Max)	1.20 %	Copper	(Max)	33 PPM
Phosphorus	(Min)	0.35 %	Selenium	(Min)	0.2 PPM
			Vitamin A	(Min)	12,000 IU/LB

INGREDIENTS

Grain Products, Forage Products, Processed Grain Byproducts, Roughage Products (15%), Soybean Meal, Molasses Products, Ground Limestone, Soybean Oil, Ammonium Chloride, Sodium Bicarbonate, Salt, Animal Protein Products, Yeast Culture, Zinc Proteininate, Manganese Proteininate, Copper Proteininate, Dried Yeast Fermentation Solubles, Cobalt Proteininate, Brewers Dried Yeast, Dried *Saccharomyces Cerevisiae* Fermentation Solubles, Hydrated Sodium Calcium Aluminosilicate, Silicon Dioxide, *Yucca Schidigera* Extract, Vitamin E Supplement, Calcium Carbonate, Mineral Oil, Thiamine Mononitrate, Vitamin A Supplement, Vitamin D3 Supplement, Magnesium Oxide, Selenium Yeast, Cobalt Carbonate, Copper Chloride, Copper Sulfate, Ethylenediamine Dihydriodide, Manganese Sulfate, Manganous Oxide, Zinc Sulfate and Zinc Oxide.

FEEDING DIRECTIONS

Feed PRODUCERS SHOW GOAT at the rate of 2.5 to 5 pounds per head per day. Provide adequate amounts of forage (hay or pasture), with unlimited access to fresh, clean water. Adding grain products to goat diet can often result in the development of urinary calculi. Consult your nutritionist before making such modifications.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats.

Manufactured by
PRODUCERS COOPERATIVE ASSOCIATION
Bryan, Texas 77806

Net Weight 50 lbs. (22.68 kg) or Bulk