



Purina® Wind and Rain® All Season 4 Mineral Tub Complete Added Zinc and Copper

Supplement feed for cattle on pasture

Weight 225lbs

GUARANTEED ANALYSIS

Crude Protein, min	5.00%	
Crude Fat, min.....	1.00%	
Crude Fiber, max.....	2.00%	
Calcium (Ca), min	4.00%	Max.....5.00%
Phosphorus (P), min.....	4.00%	
Salt (NaCl). Min	9.00%	Max.....10.80%
Magnesium (Mg), min.....	1.00%	
Potassium (K), min	2.00%	
Manganese (Mn), min.....	1,250 ppm	
Cobalt (Co), min	30 ppm	
Copper (Cu), min.....	1,250 ppm	
Selenium (Se), min	10 ppm	
Zinc (Zn), min.....	3,750 ppm	
Vitamin A, min.....	80,000 IU/Lb	
Vitamin D ₃ , min	20,000 IU/Lb	
Vitamin E, min.....	100 IU/Lb	

INGREDIENTS

Molasses Products, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Processed Grain By-Products, Calcium Carbonate, Hydrolyzed Vegetable Oil, Magnesium Oxide, Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Vitamin A Supplement, Vitamin D₃ Supplement, Vitamin E Supplement.

FEEDING DIRECTIONS

1. Feed one tub per 25-30 head of cattle and at least 2 tubs per pasture to ensure adequate intake by younger, timid animals
2. Place tubs within 100 feet of where animals congregate (loafing, grazing, feeding and watering areas)
3. Feed tubs continuously along with plentiful sources of forage and fresh, clean water
4. Cattle will consume approximately ¼ to ½ pound of this supplement per head daily.
5. If additional mineral and vitamin fortification is needed, provide a nutritionally balanced mineral/vitamin supplement containing salt.
6. This product does not contain added salt; therefore supplemental salt should be offered at all times.

Placement of tubs can influence grazing distribution in large pasture situations. Place tubs in areas on under-utilization in order to entice cattle to specific locations within a pasture.

During the first week of use, place the tubs within 100 feet of the loafing and/or watering areas. Also consider animal condition, forage quality and quantity and seasonal weather conditions. If over-consumption occurs for more than two weeks despite tub repositioning, remove tubs and evaluate the overall feeding program.

Caution: Do not feed to sheep or goats. Use only as directed.