

PRODUCERS

85 GOAT PELLET DX

MEDICATED

For Growing, Breeding and Lactating Goats whose milk will not be collected for human consumption.

(FOR RUMINANTS ONLY)

For the prevention of coccidiosis in young goats caused by *Eimeria cristenseni* and *Eimeria ninakohlyakimovae*. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

ACTIVE DRUG INGREDIENT

Decoquinatate 27.24 g/ton

GUARANTEED ANALYSIS

Crude Protein*	(Min)	15.00 %			
(Includes not more than 0.9% equivalent protein from non-protein nitrogen.)					
Crude Fat	(Min)	3.50 %	Salt	(Min)	0.50 %
Crude Fiber	(Max)	15.00 %	Salt	(Max)	0.70 %
Calcium	(Min)	0.80 %	Copper	(Min)	14 PPM
Calcium	(Max)	1.20 %	Copper	(Max)	19 PPM
Phosphorus	(Min)	0.40 %	Selenium	(Min)	0.3 PPM
			Vitamin A	(Min)	15,000.00 IU/LB

INGREDIENTS

Grain Products, Processed Grain Byproducts, Roughage Products (20%), Forage Products, Soybean Meal, Molasses Products, Ground Limestone, Ammonium Chloride, Salt, Vitamin A Supplement, Zinc Sulfate, Manganese Sulfate, Manganous Oxide, Zinc Oxide, Calcium Carbonate, Sodium Selenite, Sodium Molybdate, Ethylenediamine Dihydriodide, Cobalt Carbonate, Vitamin E Supplement, Vitamin D3 Supplement, Mineral Oil and Copper Sulfate.

FEEDING DIRECTIONS

Feed GOAT PELLET DX for at least 28 days during periods of exposure, or when experience indicates that coccidiosis is likely to be a hazard. Provide adequate amounts of forage (hay or pasture), with unlimited access to fresh, clean water. Adding grain products to goat diets can often result in the development of urinary calculi. Consult your nutritionist before making such modifications.

<u>Goat Wt. in Lbs</u>	<u>Lbs/Day Goat Pellet DX</u>
20	0.33
40	0.67
60	1.00
80	1.34

WARNING: Do not feed to goats producing milk for food.

Manufactured by
PRODUCERS COOPERATIVE ASSOCIATION
Bryan, Texas 77806

Net Weight 50 lbs. (22.68) or Bulk