

# PRODUCERS

## 85/2062 GOAT PELLET DX

### MEDICATED

**FOR GROWING, BREEDING AND LACTATING GOATS WHOSE MILK WILL NOT BE COLLECTED FOR HUMAN CONSUMPTION (FOR RUMINANTS ONLY)**

For the prevention of coccidiosis in young goats caused by *Eimeria christenseni* and *Eimeria ninakohlyakimovae*. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

#### ACTIVE DRUG INGREDIENT

Decoquinatate ..... 27.2 g/ton

#### GUARANTEED ANALYSIS

Crude Protein	(Min)	15.00 %			
Includes not more than 1.20% equivalent non-protein nitrogen.					
Crude Fat	(Min)	3.00 %	Salt	(Min)	0.40 %
Crude Fiber	(Max)	13.00 %	Salt	(Max)	0.70 %
Calcium	(Min)	0.90 %	Copper	(Min)	19 PPM
Calcium	(Max)	1.30 %	Copper	(Max)	21 PPM
Phosphorus	(Min)	0.35 %	Selenium	(Min)	0.3 PPM
			Vitamin A	(Min)	15,000 IU/LB

#### INGREDIENTS

Grain Products, Roughage Products (20%), Processed Grain Byproducts, Forage Products, Soybean Meal, Molasses Products, Ground Limestone, Ammonium Chloride, Salt, Soybean Oil, Calcium Carbonate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Selenium Yeast, Cobalt Carbonate, Copper Chloride, Copper Sulfate, Ethylenediamine Dihydriodide, Manganese Sulfate, Manganous Oxide, Zinc Sulfate, Zinc Oxide and Sodium Selenite.

#### FEEDING DIRECTIONS

Feed GOAT PELLET DX for at least 28 days during periods of exposure, or when experience indicates that coccidiosis is likely to be a hazard. Provide adequate amounts of forage (hay or pasture), with unlimited access to fresh, clean water. Adding grain products to goat diets can often result in the development of urinary calculi. Consult your nutritionist before making such modifications.

<u>Goat Wt. in Lbs</u>	<u>Lbs/Dav Goat Pellet DX</u>
20	0.33
40	0.67
60	1.00
80	1.34

WARNING: Do not feed to goats producing milk for food.

Manufactured by  
PRODUCERS COOPERATIVE ASSOCIATION  
Bryan, Texas 77806

Net Weight 50 lbs. (22.68 kg) or Bulk