

PRODUCERS

91/1024

STOCKER STARTER DX-1.5

MEDICATED
FOR CATTLE ON PASTURE

For the prevention of coccidiosis in ruminating and non-ruminating calves and cattle caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT

Decoquinatate 30.20 g/ton (0.0033 %)

GUARANTEED ANALYSIS

Crude Protein	(Min)	12.00 %	Phosphorus	(Min)	0.40 %
Crude Fat	(Min)	2.90 %	Salt	(Min)	0.45 %
Crude Fiber	(Max)	21.70 %	Salt	(Max)	0.85 %
Calcium	(Min)	0.60 %	Potassium	(Min)	1.20 %
Calcium	(Max)	1.00 %	Vitamin A	(Min)	8,000 IU/LB

INGREDIENTS

Roughage Products (34%), Forage Products, Processed Grain Byproducts, Grain Products, Roughage Products, Plant Protein Products, Molasses Products, Ground Limestone, Calcium Lignosulfonate, Salt, Selenium Yeast, Cobalt Carbonate, Copper Chloride, Copper Sulfate, Ethylenediamine Dihydriodide, Manganese Sulfate, Manganous Oxide, Zinc Sulfate, Zinc Oxide, Mineral Oil, Calcium Carbonate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Calcium Propionate, Cottonseed Meal, Yeast Culture, Vitamin D3 Supplement, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex and Cobalt Glucoheptonate.

FEEDING DIRECTIONS

Feed at a rate of 1.5 pounds per 100 pounds bodyweight per day to provide 22.7 mg of decoquinatate per 100 pounds bodyweight per day. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Always provide adequate amounts of good forage (hay or pasture) and plenty of fresh, clean water.

WARNING: Do not feed to cows producing milk for food. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Feed type C feed or milk replacer to provide 22.7 mg per 100 lb. bodyweight (0.5 mg/kg) per day.

Manufactured by
PRODUCERS COOPERATIVE ASSOCIATION
Bryan, Texas 77806

Net Weight 50 lbs. (22.68 kg) or Bulk

1024 6 7/11/2020

MFG: ~DATE

~MTIME

