



# Purina® Catfish 32

**Specie:** Fish & Aquatics **Type of Feed:** Complete **Form of Feed:** Extruded

## General Description:

A floating fish feed that is comprised of high-quality ingredients and nutrient levels designed to maximize efficient production of all types of catfish. When combined with proper management practices, Purina® Catfish 32 feed is formulated to support the growth of large, healthy catfish.

**PURINA®**  
**CATFISH 32**  
FEED FOR CATFISH

**CAUTION: FEED AS DIRECTED**

## GUARANTEED ANALYSIS

**Crude Protein, (Min)**.....**32.00%**  
**Crude Fat, (Min)**.....**4.00%**  
**Crude Fiber, (Max)**.....**7.00%**  
**Phosphorus (P), (Min)**.....**0.80%**  
**Ascorbic Acid, (Min)**..... **90 mg/kg**

## \*INGREDIENTS

Dehulled Soybean Meal, Wheat Middlings, Ground Corn, Poultry By-Product Meal, Calcium Carbonate, Fish Oil Preserved with Ethoxyquin, Animal Fat Preserved with BHA and BHT, Fish Meal, Dicalcium Phosphate, DL-Methionine, L-Ascorbyl-2-Polyphosphate (Stabilized Vitamin C), Choline Chloride, Vitamin E Supplement, Niacin Supplement, d-Calcium Pantothenate, Riboflavin Supplement, Thiamine Mononitrate, Biotin, Pyridoxine Hydrochloride, Folic Acid, Vitamin A Supplement, Vitamin D3 Supplement, Menadione Sodium Bisulfate Complex, Vitamin B12 Supplement, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Sodium Selenite, Cobalt Carbonate, Ethylenediamine Dihydroiodide, Ethoxyquin (a Preservative).

**\*Note: Ingredients differ by manufacturing plant.**  
**Please refer to the tag for a current list of ingredients.**

## FEEDING DIRECTIONS:

1. Feed Purina® Catfish 32 feed to pond culture catfish or other farm pond fish 6 to 7 inches in size or larger when stocking rates are 4000 or less fish per acre.
2. Ensure that adequate oxygen is available to the fish and follow the recommendations for various water temperatures.

3. Use the following guide when adequate oxygen is available to the fish:

Water Temperature	Amount to feed daily	Frequency of feeding
65-75° F	2-3% of body weight	Daily 6 days/wk.
75-85° F	3-5% of body weight	Daily 6-7 days/wk
85-90 °F	2-3% of body weight	Daily 5-6 days/wk

4. For best results, feed as much Catfish 32 feed as the fish will consume in 15-20 minutes.
5. If feed is completely consumed within 15 minutes, gradually offer the fish additional feed.
6. If feed is still floating on the surface after 20 minutes, the quantity of feed offered should be decreased.

## LIMITATIONS FOR USE:

**Available Options:**

Product No.	Package	Size Diameter
0040967	Bag 50#	5/16 inch

**Product Features:**

32% High-quality Protein

High Energy Content

Stabilized Vitamin C

**Product Benefits:**

Balanced amino acids support lean growth.

Encourages rapid and efficient growth.

Supports optimal health.

**IMPORTANT:**

This product is to be fed only to the animal species as directed on this label. Follow these management practices:

1. Poor water quality is the most common cause of fish loss.
2. Avoid overstocking ponds; exceeding the biomass capacity of a pond may result in fish loss.
3. Avoid overfeeding; excessive nutrient load will deplete water oxygen.
4. Reduce feeding frequency to not more than five days per week when water temperature reaches 85° F. Drastically reduce feeding rates and reduce frequency of feeding to three days per week at temperatures above 95° F.
5. Keep ponds adequately aerated.
6. If feed is not completely eaten within 20 minutes, reduce the amount that is offered.
7. Feed in late afternoon/early evening, when dissolved oxygen concentration is highest.

**CAUTION:**

Always follow label directions.

Store in a dry, well-ventilated area protected from rodents and insects.

Do not feed moldy or insect-infested feed to animals as it may cause illness or death.

Please consult the tag sewn onto the bag for complete feeding instructions.

**KEY POINTS:**

1. A feeding program is only as effective as the management practices followed.
2. Consumption can vary depending upon water temperature, season of the year, nutritional needs of the fish, availability of other foods and fish familiarity with particle feed.